



TOT WATCH



**MENINGITIS &
SEPTICAEMIA**
**Higher risk
for young
children**

What every parent & carer should know

If you have a child under five years old, you need to be aware of meningitis and septicaemia because children in this age group are more at risk than older children and adults. Meningococcal disease is the leading cause of meningitis amongst children in the UK and Ireland. This bug kills more children under five than any other infectious disease. Even children who survive may be left with life-changing disabilities from learning difficulties and behavioural problems to hearing loss and amputations.

Fortunately, most children have natural resistance to the bugs that cause meningitis and septicaemia and vaccines give excellent protection against many forms. But not all forms can be prevented, so it's very important that everyone is aware of the signs and symptoms.

Meningitis and septicaemia can be hard to recognise, especially in small children who get lots of minor illnesses with similar symptoms and can't explain how they are feeling.

But these diseases can kill in hours so trust your instincts and seek urgent medical help if you suspect meningitis or septicaemia.

What are meningitis & septicaemia?

Meningitis is inflammation of the lining around the brain and spinal cord - the meninges. Septicaemia is blood poisoning, and is the more life-threatening form of the disease.



Symptoms to watch out for

The first symptoms are usually fever, vomiting, headache and feeling unwell. **Red ticks show symptoms that are more specific to meningitis and septicaemia and less common in milder illnesses.** Limb pain, pale skin and cold hands and feet often appear earlier than the rash, neck stiffness, dislike of bright lights and confusion.

	Septicaemia	Meningitis
Fever and/or vomiting 	✓	✓
Severe headache 		✓
Limb/joint/muscle pain <small>(sometimes stomach pain/diarrhoea)</small> 	✓	
Cold hands and feet/shivering 	✓	
Pale or mottled skin 	✓	
Breathing fast/breathless 	✓	
Rash <small>(anywhere on the body)</small> 	✓	✓
Stiff neck <small>(less common in young children)</small> 		✓
Dislike of bright lights <small>(less common in young children)</small> 		✓
Very sleepy/vacant/difficult to wake 	✓	✓
Confused/delirious 	✓	✓
Seizures (fits) may also be seen 		✓

Other signs in babies

- Unusual grunting sounds
- Tense or bulging soft spot on their head
- Refusing to feed
- Irritable when picked up, with a high pitched or moaning cry
- A stiff body with jerky movements, or else floppy and lifeless
- Fever is often absent in babies less than three months of age

The tumbler test

If you are seriously worried about a child who is ill, **don't wait for a rash to appear – get medical help**. But if your child is ill and gets a new rash or spots, use the Tumbler Test.

Press a clear glass tumbler firmly against the rash. If you can see the marks clearly through the glass **seek urgent medical help immediately**.

Check the entire body. Look out for tiny red or brown pin-prick marks which can change into larger red or purple blotches and blood blisters.

The darker the skin the harder it is to see a septicaemic rash so check lighter areas like palms of hands and soles of the feet or look inside the eyelids and the roof of the mouth.



Things to remember

- Don't wait for a rash as it doesn't always appear
- Not everyone gets all of these symptoms
- Symptoms can appear in any order
- Septicaemia can occur with or without meningitis
- You know your child best; check on them often, trust your instincts and act fast.

How can I protect my children from meningitis?

The single most effective thing you can do to protect your children from meningitis is to make sure they are fully immunised. Every injection in the routine immunisation programme for babies protects against some form of meningitis. Booster vaccines are not just a top-up. Vaccines given after 12 months of age are very important. Without these immunisations your child's protection will be short lived.

But since not all forms of meningitis can be prevented, it's important you know the symptoms. That way you can recognise the disease in time to get medical help if your child is affected.

Trust your instincts

You know your child best - you are with them every day. You are the best person to spot the difference in behaviour or in the sound of a cry. Remember, meningitis and septicaemia strike fast. If you think your child is seriously ill get medical help immediately. Children with meningitis or septicaemia will usually get ill quickly and get worse fast, so check them often, including during the night.

About us



Our vision is for a world free from meningitis and septicaemia. That's why we fund research into the prevention, detection and treatment of the diseases, promote education and awareness amongst the public and health professionals and provide support to those affected.

Our Patron, Paralympic gold medallist Jonnie Peacock with Junior Ambassador Sofia Crockatt and her running coach Vicky Huyton

For information and support our Freephone helpline is available 365 days a year

080 8800 3344 (UK)
1800 41 33 44 (Ireland)
helpline@meningitis.org
www.meningitis.org



We need your help

We are committed to saving lives and have invested over £18m/€24m in vital vaccine research and studies which speed up diagnosis and improve treatment. We also spend around £1m/€1.2m a year supporting families and raising awareness of the diseases among the public and health professionals with free information. If this leaflet has inspired you to help by donating or fundraising, please contact your local office or visit us online.

Our offices

Bristol

Tel 0333 405 6262
info@meningitis.org

Edinburgh

Tel 0131 510 2345
info@scotland-meningitis.org.uk

Belfast

Tel 028 9032 1283
info@meningitis-ni.org

Dublin

Tel 01 819 6931
info@meningitis-ireland.org

A charity registered in England and Wales no 1091105, in Scotland no SC037586 and in Ireland 20034368

Registered Office: Newminster House,
Baldwin Street, Bristol BS1 1LT

© Meningitis Research Foundation 01/2016

